



BULLETIN 2

This second Bulletin expands on the priority groups requiring a second dose of **CELVAPAN** as agreed at the National Public Health Emergency Team meeting on 12th November

Everyone vaccinated with **CELVAPAN** requires two doses at least three weeks apart

At the mass vaccination clinics priority for the 2nd dose of Celvapan should be given to

- **at risk children aged 6 months to 12 years**
- **those 13 years and over who are immunocompromised**
- **pregnant women**
- **all those with cystic fibrosis**

How to identify this high priority group?

Please contact Robert Cooke, PCCC I.C.T. Programme Manager at robert.cooke@hse.ie if you wish to receive a daily list by clinic of those in these at risk groups who have received their first dose of vaccine.

NOTE: This will only be possible where the details of the “at risk” group have been completed so it is important to ensure this information is collected at the first visit

It will not be possible to identify those with cystic fibrosis as they will have been coded as “respiratory disease”. The Cystic Fibrosis Association will contact them directly to attend the same MVC

All those in the high priority groups above should be contacted to attend for their 2nd dose after three weeks.

REMEMBER you will need to make sure you have supplies of CELVAPAN to give this group their second dose of vaccine.

All others who receive Celvapan should be told that they may need a second dose of vaccine at a later date in which case they will be contacted